

		Bedarf an kg Stickstoff (N) / ha														Bed. In Gramm / 10 m ²																							
		<div style="display: flex; justify-content: space-between; font-size: small;"> <5510152025303540455055>60 </div>																																					
		Tage														Monate																							
		Wochen														Ca K P2O5 N																							
Kulturperiode		1	2	3	4	5	6	7	8	9	10	11	12	13	14	3	4	5	6	7	8	9	10	11	12	13	14	15	16	5	6	7	8	9	Ca	K	P2O5	N	
Artischocke	FB; gpf															E																							
Brokkoli	FB; gpf																																						
Brokkoli																																							
Bohne	FB; gpf																																						
Chinakohl	FB; gs																																						
Dill	FB; gs																																						
Gurke	FB; gs																																						
Gurke Salat	KK; gpf																																						
Salat (Eissalat)	FB;gpf																																						
Salat (Endivie)	FB;gpf																																						
Erbsen	FB; gs																																						
Kohl	FB; gpf																																						
Kohl (Vlies)	FB; gpf																																						
Kohlrabi	FB; gpf																																						
Karotte	FB; gs																																						
Kochsalat	FB; gpf																																						
Kraut (Vlies)	FB; gpf																																						
Krenn	FB; gpf																																						
Kürbis	FB; gpf																																						
Mangold	FB; gpf																																						
Melanzani	FB; gpf																																						
Melanzani	GH; gpf																																						
Paprika	FB; gpf																																						
Petersilie	FB; gs																																						
Radieschen	FB; gs																																						
Radieschen	GH; gs																																						
Rhabarber	FB; gpf																																						
Schnittlauch	FB; gs																																						
Tomaten	GH; gpf																																						
Mais (Zucker)	FB; gpf																																						
Erdbereren																																							
Weinreben																																							
Hochstammobst																																							

Guter Stallmistdüngung senkt den Bedarf um 30 %